GENERAL INDEX TO VOLUME X

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italic*; and previously published scientific names and all other matter, in ordinary type.

Acidity, relation of, to growth of citrus-destroying fungi, 245; to Diaporthe Sojae, 157 Actinastrum Hantzschii, 416 Aetheolaena involucrata, 83 Agaricus, 181
Alcoholic products, production of, by various fungi, 290 Aleurodiscus perideniae, 187 Algae from lakes in the northeastern part of North Dakota, 393 Alternaria Citri, 242, 281 Amino nitrogen, determination of, 330 Amoeba or protozoan theory of mosaic disease, 196 Anabaena affinis, 403; circinalis, 404; floe-aquae, 404, var. Treleasii, 404; Lemmermanni, 404; macrospora, 404, 422; spiroides, 405, 422 Ankistrodesmus falcatus, 415; spiralis, 415

Apparatus and materials used in sugar determination, 72 applanatus (Fomes), 184
arachnoideum (Corticium), 187
arcularius (Polyporus), 182
Armstrong, Joanne K., B. M. Duggar
and. Indications respecting the nature
of the infective particles in the mosaic
disease of tobacca 101

Aphanizomenon flos-aquae, 405 Aphanochaete repens, 419

disease of tobacco, 191 Arthrodesmus controversus, 410 Arthrospira Jenneri, 402

Asci of soybean fungus, 124, containing ascospores, 122, germinating, 124
Aspergillus niger, 242; nitrogen metabolism in, 322
Astronousiis generaline, 408, 422

Asterocystis smaragdina, 406, 422 auricula-Judae (Auricularia), 188 Auricularia auricula-Judae, 188; nigres cens, 188; tenuis, 188 australis (Fomes), 184

B

Bacillus atrosepticus, 5, 17, 43; caulivorus, 3; coli, 46; communis, 46; fluorescence liquefaciens, 3; melanogenes, 19; phytophthorus, 5, 17; solanisaprus, 18; vulgatus, 46
Bacterial theory of mosaic disease, 194
Beals, Cora Mauts. An histological study of regenerative phenomena in plants, 369 study of regenerative phenomena in plants, 369
Blackleg, potato, 1; diagnosis of, 9; economic aspects of, 13
Botryococcus Braunii, 413
Bryophyllum, experiments in regeneration with, 373, 378, 384
Bulbochaete sp., 417
Burt, E. A. Higher fungi of the Hawaiian Islands, 179

Cacalia arborea, 77; dichroa, 87; glabrata, 97; involucrata, 83; patens, 87; pul-chella, 90; teretifolia, 96; vaccinioides, 97

Callothrix fusca, 405; scytonemicola, 405 Canp, A. F. Citric acid as a source of carbon for certain citrus fruit-destroying fungi, 213 Campanella cucullata, 186 Carbohydrate utilization by strains of the blackleg bacillus and other microorganisms, 45

organisms, 45 Carbon: analysis apparatus, 237, 238, 298; citric acid as a source of, for certain fruit-destroying fungi, 213; total, determination of, 236
Carter, Nellie, George T. Moore and. Algae from lakes in the northeastern

part of North Dakota, 393 cepaeforme (Lycoperdon), 189 Chaetophora elegans, 418

(423)

ANN. Mo. Bor. GARD., Vol. 10, 1923

Chara elegans, 420
chioneus (Polyporus), 182
Chondrocystis Schauinslandii, 398
Chroococcus dispersus, 397, var. minor, 397; limneticus, 397; turgidus, 397
cinnamomea (Hymenochaete), 186
Citrate, growth of fungi on varying percentages of, 255
Citric acid, 213; apparatus, 222; as a source of carbon for certain citrus fruit-destroying fungi, 213; discussion of, 214; occurrence of, 226; physiological role of, 229; qualitative detection of, 215; quantitative determination of, 216
Cladophora Kuetsingiana, 419
Clathrocystis aeruginosa, 399, var. major, 399; holsatica, 399
Closterium Dianae, 407, var. arcuatum, Chara elegans, 420 Closterium Dianae, 407, var. arcuatum, 408; eboracense, 408, 422; lanceolatum, 408
Coelastrum cambricum, 415; microtum, 408
Coelastrum cambricum, 415; microporum, 415
Coelosphaerium Kuetzingianum, 400;
Naegelianum, 400
coliforme (Myriostoma), 189
coliformis (Geaster), 189
commune (Schisophyllum), 181
Conferva bombycina f. tenuis, 407
Corium (Myreonastrum), 189 Corium (Myconastrum), 189
Corium (Myconastrum), 189
corrugata (Trametes), 185
Corticium arachnoideum, 187; granu-Cosmarium calcareum, 409; formosulum, 409, var. Nathorstii, 409; granatum, 409; hians, 409; impressulum, 409; Meneghinii, 409; pygmaeum, 409; Regnellii, 409; scopulorum, 409; subcostatum f. minor, 409; tenue, 409; Turpinii var. podolicum, 409
Crepidotus fulvotomentosus, 181; rhizomorphys. 181 morphus, 181 Crucigenia quadrata, 414; tetrapedia, 414 cucullata (Campanella), 186 cucullata (Laschia), 186 cucullatus (Merulius), 186 cucullatus (Merulius), 186 Cylindrocapsa geminella, 417 Cyphella villosa, 186

D

Daedalea sanguinea, 185
Dextrose solution, 236, 323
Diaporthe Sojae, 128, 170-178
Dictyosphaerium Ehrenbergianum, 415;
pulchellum, 415
Diplodia natalensis, 242; nitrogen metabolism in, 322
dryophilus (Polyporus), 182

Duggar, B. M., and Joanne Karrer Armstrong. Indications respecting the nature of the infective particles in the mosaic disease of tobacco, 191 Duggar's solution, 323

E

elegans (Stereum), 187
Enteromorpha intestinalis, 417; prolifera, 417
Enzyme theory of mosaic disease, 194
Epithele hydnoides, 188
Eudorina elegans, 412
Exidia polytricha, 188

F

fasciatus (Fomes), 184
fasciculata (Poria), 184
fasciculatus (Fomes), 183
fibula (Polystictus), 184
fabellaris (Polyporus), 182
fabellatus (Pieurotus), 180
Flagellata, 397
Flax, experiments on regeneration with, 374, 378, 380
floccosus (Polystictus), 184
Fomes applanatus, 184; australis, 184; fasciatus, 184; fasciatus, 184; fasciatus, 183; Fullageri, 184; tawaiensis, 183; Korthalsii, 182; rimosus, 184; robustus, 184; senez, 182
Fullageri (Fomes), 184
fulvotomentosus (Crepidotus), 181
Fungi: certain citrus fruit-destroying, citric acid as a source of carbon for, 213; growth of, on varying percentages of citrate, 255; some aspects of nitrogen metabolism in, 299; studies in the physiology of the, XVI, 299

G

Ganoderma, 184
Geaster coliformis, 189
gemmatum (Lycoperdon), 189
gilvus (Polyporus), 182
glareosa (Psathyra), 181
Gloeocapsa fenestralis, 397
Gloeotrichia echinulata, 406
Glucose solution, 323
Gomphosphaeria aponina, 400, var.
cordiformis, 400
granulare (Corticium), 187
Greenman, J. M. Studies of South
American Senecios—I, 73
Griading, effect of, on the infectivity of
tobacco virus, 204

Gynoxis alternifolia, 76

H

hawaiensis (Fomes), 183 Hawaiian Islands, higher fungi of the,

Herposteiron confervicola, 419 Higher fungi of the Hawaiian Islands,

179 Hirneola nigra, 188 hirsutus (Polystictus), 184 Histological study of regenerative phenomena in plants, 369 Holopedia irregularis, 400

Horseradish, experiments on regenera-tion of roots of, 371, 578, 382 hydnoides (Epithele), 180

Hydnum, 186 Hydrogen-ion concentration of culture solutions, 325; change of media, 335; effect of, on germination of pycnospores of Diaporthe Sojae, 153; of spores of Diaporthe Sojae, 153; of culture solutions after growth of fungous mats, 258; of citric acid solution after fermentation by various fungi, 260; of dextrose solution after fermentation by various fungi, 259; reaction of, to growth of citrus fruit-destruing funcionals.

destroying fungi, 245 Hymenochaete cinnamomea, 186; spreta, 186; tenuissima, 186

I

Indications respecting the nature of the infective particles in the mosaic disease of tobacco, 191

Jennison, H. M. Potato blackleg, with special reference to the etiological agent, 1

K

Kirchneriella lunaris var. irregularis, 415; obess, 415, var. aperta, 415 Klotz, L. J. Studies in the physiology of the fungi, XVI, Some aspects of nitrogen metabolism in fungi, 299 Korthalsii (Fomes), 182

L

lactinea (Trametes), 185 Laschia cucullata, 186 latum (Stereum), 187

Lehman, S. G. Pod and stem blight of soybean, 111 Lemons, acid and sugar content of, 227 Lepiota xylophila, 180
Light, effect of, on production of pycnidia of various fungi, 147 lignosus (Polyporus), 182 Lycoperdon cepaeforme, 189; gemma-tum, 189; Wrightii, 189 Lyngbya birgei, 402; contorta, 403; Martensiana, 403

M

marginata (Pholiota), 181 Media: employed in study of blackleg organism, 26; used in culture of soyorganism, 26; used in culture of soy-bean fungus, 153; used in nitrogen experiments, 323 Melilotus alba, 120 Merismopedia glauca, 400; irregularis, 400; punctata, 400; tenuissima, 400 Merulius cucullatus, 186 Microchaete corymboso, 86; glabrata, 97; pulchella, 90; trichopus, 90; vaccinioides, 97 Micrococcus phytophthorus, 2 Microcystis incerta, 399; pulvera, 399; stagnalis, 399 microloma (Polystictus), 184 Moisture, effect of, on pycnidia forma-Mosture, energy, on pythalic tion, 152

Moore, George T., and Nellie Carter.

Algae from lakes in the northeastern
part of North Dakota, 393 Mosaic disease of tobacco, indications respecting the nature of the infective particles in the, 191 Mougiotis calcarea, 412 Mycenastrum Corium, 189

N

Myriostoma coliforme, 189

Naucoria triscopoda, 181 Nectria Ipomoeae, nitrogen fixation of, 334 nigra (Hirneola), 188
nigrescens (Auricularia), 188
Nitrites, determination of, 329
Nitrogen metabolism in fungi, some Nitrogen metabolism in rungi, some aspects of, 299
Nitrogen, total, determination of, 327
Nodularia spumigena, 403, var. genuina, 403, var. major, 403
North Dakota: algae from lakes in the northeastern part of, 393; map of the physiographic regions of, 390; map of northeastern part of, 392; notes on the physiography of, and the conditions in certain of its waters, 383

Odontia Wrightii, 186 Oedogonium sp., 417 Oocystis crassa, 413: lacustris, 413: pusilla, 413 Oospora Citri-auranti, 244 Oranges, acid and sugar content of, 227 Oscillatoria amphibia, 401; brevis, 401; chalybea, 401; janthiphora, 401, 422 ostreatus (Pleurotus), 180 Oxalic acid, determination of, 236

P Pandorina morum, 412 Pediastrum boryanum, 416, var. longi-corne, 416; duplex var. clathratum, 416, var. gracillimum, 416; simplex, 417, var. duodenarium, 417; tetras, 417 Penicillium digitatum, 241; italicum, 240; stoloniferum, 241 Peptone solutions, 323 Periode solutions, 225 Peronospora infestans, 2 Persoonii (Polystictus), 185 Pholiota marginata, 181 Phoma Betae, 322; nitrogen metabolism in, 334 Phoma blight of soybean, 111 Phomopsis Citri, 244, 289 Physiography of North Dakota, notes on, and the conditions in certain of its and the conditions in certain of its waters, 385 pinsitus (Polystictus), 184 Pleurotus flabellatus, 180; ostreatus, 180 Pod and stem blight of soybean, 111, 170–178; description of, 112; morphology of, 115

Polycystis pallida, 399; stagnalis, 399

Polyporus arcularius, 182; chioneus, 182; dryophilus, 182; flabellaris, 182; gilvus, 182; lignosus, 182; sulphureus, 182 182 Polystictus fibula, 184; floccosus, 184; hirsutus, 184; microloma, 184; Per-soonii, 185; pinsitus, 184 polytricha (Exidia), 188 Poria fasciculata, 184; sp., 185 Potato blackleg, with special reference to the etiological agent, 1 Potato plants, affected with blackleg, showing characteristic symptoms, 70. showing characteristic symptoms, 70, 72 Protozoan theory of mosaic disease, 197 Peacalium vaccinioides, 97; glabratum, 97

Psathyra, 181; glareosa, 181 Pycnospores of soybean fungus, 119; germinating, 120 Pycnosporophores of soybean fungus,

R

Rainfall, relation of amount of, to prevalence of pod and stem blight of soybean, 145
Regenerative phenomena in plants, an histological study of, 369
Rhabdoderma sigmoidea, 398, 422, f. minor, 398, 422
Rhiscolonium crispum, 410: hisroglyphia Rhizoclonium crispum, 419; hieroglyphicum, 419 rimosus (Fomes), 184 Rivularia coadunata, 406 robustus (Fomes), 184

sanguinea (Daedalea), 185
Seenedesmus bijugatus, 413, var. alternans, 413, var. flexuosus, 413; dimorphus, 414; obliquus, 414; quadricauda, 414, var. bicaudatus, 414, 422, var. quadrispina, 414, var. Westii, 414
Schizophyllum commune, 181
Sclerotinia Libertiana, 244, 287
Selenastrum gracile, 415; Westii, 415
Senecio aberrans, 73, 100; abietinus, 75; adenophyllus, 76; andicola, 76; apiculatus, 76; arboreus, 77; aridus, 77, 102; attenuatus var. microphyllus, 80; auritus, 78; Chinogeton f. macrocephalus, 80; chaquiroensis 79; clavifolius, 80; comosus, 80; culcitioides, 81; decompositus, 81; ericaefolius, 81; formosus, 82; formosus, 93, var. subruncinatus, 94; genuficaus, 82; involucratus, 83; iodopappus, 83; laciniatus, 84; latiforus, 84; ledifolius, 76; lophophilus, 84, 104; Magnusii, 85; medullosus, 85; microchaete, 86; modestus, 86; nubigenus, 87; organensis, 87; otophorus, 87; patens, 87; pellucidinervis, 88; Pennellif, 88, 106, var. gachetensis, 89; pericaulis, 89; pimpinellifolius var. laciniata, 84, var. nubigena, 87; pindilicensis, 90; pulchellus, 90; rudbeckiaefolius, 90; sanctae-martae, 91; acortifolius, 91; sinapoides, 92; Sprucei, 93; stigophlebius, 93; stuebelii, 93; subdecurrens, 93; subglomerosus 93, 108; sanguinea (Daedalea), 185

subruncinatus, 94; sylvicolus, 95, 110; Tabacon, 82; teretifolius, 96; theae-folius, 97; tolimensis, 97; vaccinioides, 97, var. pruinosus, 97; vernicosus var. major, 76, var. microphyllus, 97; Weddellianus, 97; Xanthopappus, 98
Senecios, studies of South American—I, 72

senex (Fomes), 182 Soja max, 129 Solanum tuberosum, 1 Sorastrum spinulosum, 415

Soybean fungus, 170-178
Soybean, pod and stem blight of, 111;
cultural characters, 142; manner of
infection, 139; overwintering and dissemination, 140; varietal susceptibility,

Sphaeropsis malorum, change in fungous mat, 340; nitrogen metabolism in fungi, 322

fungi, 322
Spirogyra lutetiana, 412
Spirulina major, 402; subtillissima, 402; tenerrima, 402
spreta (Hymenochaete), 186
Staurastrum gracile, 410, forma, 410, 422; paradoxum var. evolutum, 411, 422; f. biradiata, 411, 422; tetracerum f. evolutum, 411
Stereum elegans, 187: latum, 187

Stereum elegans, 187; latum, 187 Stigeoclonium sp., 418, 422 Stylospores of soybean fungus, 121 Sugars, reducing, determination of, 48, 236, 325

Sweet-potato, experiments on regenera-tion with, 371, 378, 382

tenuis (Auricularia), 189 tenuissima (Hymenochaete), 186 Tetraedron limneticum, 413; quadri-cuspidatum, 413, 422; trigonum var. respidatum, 413, 422; trigonum var. gracile, 413
Tetrapedia Reinschiana, 401
Tobacco, mosaic disease of, 191
Trametes corrugata, 185; lactinea, 185; sp., 185 triscopoda (Naucoria), 181

U

Ultrafiltration experiments in mosaic disease, 199 Ustulina vulgaris, 189

villosa (Cyphella), 186 Virus theory of mosaic disease, 195 Volvox globator, 412; mononae, 412 vulgaris (Ustulina), 189

Wollea saccata, 403 Wrightii (Lycoperdon), 189 Wrightii (Odontia), 186

xylophila (Lepiota), 180

Y

Young, R. T. Notes on the physic-graphy of North Dakota and the con-ditions in certain of its waters, 385

 \mathbf{z}

Zygnema, 411